## NONOPERATIVE ADHESIVE CAPSULITIS PT PROTOCOL

Name:	Date:
Diagnosis:	
Date of Surgery	
Frequency: 1 2 3 4 times/week	
Duration: 1 2 3 4 5 6 Weeks	

- Outpatient pain medication and modalities
  - o Ice, heat, ultrasound, etc
- Apply modalities with shoulder at end range (comfortable) position (not arm at side)
- A/AA/PROM
  - No limitations, focus on IR and ER at 90° ABD in supine position.
  - Try to preserve as much IR and ER as possible.
- Emphasize GENTLE PROM to start
- Work in pain-free arc, but emphasize modalities to stretch.
- Work on full flexion and abduction.
  - Emphasize glenohumeral motion, block scapulothoracic motion with abduction/ flexion from 0-80°.
- Rotator cuff and scapular stabilization program exercises
  - Begin at 0° and progress to 45°/90° as tolerated pain-free
- Home Exercise program
- Stretches to be done 3-4 times a day for 1-15 minutes per session

Comments:	
Functional Capacity Evaluation Work Hardening/Work Conditioni	ng Teach HEP
Modalities:	
Electric StimulationUltrasoundIontophoresisPhonophoresis TENS Heat before/after Ice before/after Trigger points mas Therapist's discretion Other	
Signature: Date:	